

Filan[®]

Filan is a protectant and systemic fungicide for use against early and late-season diseases in oilseed rape.

A water dispersible granule containing 50% w/w boscalid.

Risk and Safety Information

If medical advice is needed, have product container or label at hand.
Keep out of reach of children.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

PCS No.: 04325

To avoid risks to human health and the environment, comply with the instructions for use.

2.5 kg e

UN 3077
Packing Group III
Environmentally hazardous
substance, solid, N.O.S.
(contains boscalid 50%)
Marine Pollutant

Supplied by:
BASF Ireland Limited
P.O. Box 4, Earl Road
Cheadle Hulme, CHEADLE
Cheshire SK8 6QG, UK
Tel: 01 825 5701
Fax: 01 825 2038

Emergency Information
(24 hours freephone):
0049 180 227 3112
Technical Enquiries:
0044 (0)845 602 2553 (office hours)

Authorization holder:
BASF plc, P.O. Box 4,
Earl Road, Cheadle Hulme,
CHEADLE
Cheshire, SK8 6QG, UK


We create chemistry

® = Registered trademark of BASF

81107455IE1056



FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE, as directed below:

CROPS:	Winter and Spring Oilseed Rape
MAXIMUM INDIVIDUAL DOSE:	0.5 kg/ha
MAXIMUM TOTAL DOSE PER CROP:	1.0 kg/ha
LATEST TIMING:	Before 60% of pods have reached final size (GS 76)

READ ALL PRECAUTIONS BEFORE USE

PRECAUTIONS

WEAR SUITABLE PROTECTIVE GLOVES when handling the product.
WHEN USING DO NOT EAT, DRINK OR SMOKE.
WASH HANDS before meals and after work.

Do not contaminate water with the product or its container.
Do not clean application equipment near surface water.
Avoid contamination via drains from farmyards and roads.

STORAGE

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.
KEEP OUT OF REACH OF CHILDREN.
STORE IN ORIGINAL CONTAINER, tightly closed, in a safe place.
RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.
Store in a suitable pesticide store. Keep dry and protect from frost.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Filan is a protectant fungicide for use in oilseed rape for the control of *Sclerotinia* stem rot (*Sclerotinia sclerotiorum*) and dark leaf and pod spot (*Alternaria brassicae*). A reduction in *Phoma* leaf spot and stem canker (*Leptosphaeria maculans*) infection may also be obtained.

1. Restrictions/Warnings

For professional use only.

For use by tractor mounted/trailed sprayers only.

Avoid spray drift on to neighbouring crops.

Filan represents minimal hazard to bees when used as directed. Consideration should be given to informing bee-keepers when you intend spraying crops in flower.

A reduction in *Sclerotinia* control may be observed in oilseed rape when high-risk infection conditions occur after flowering, leading to secondary spread of the disease through plant-to-plant contact, which may be more severe in lodged crops.

2. Disease Control

Filan is a protectant fungicide for use in oilseed rape for the control of *Sclerotinia* stem rot and *Alternaria* dark leaf and pod spot. A reduction in *Phoma* leaf spot and stem canker infection may also be obtained.

2.1 Resistance

Filan contains boscalid which belongs to the SDHI (carboxamide) group of fungicides

Currently there is no evidence of field strains of *Leptosphaeria maculans*, *Sclerotinia sclerotiorum* or *Alternaria brassicae* that are resistant to the carboxamide fungicides in oilseed rape.

Currently there is evidence of cross resistance between boscalid and other fungicides in the carboxamide group, but no cross resistance has been observed with fungicides with other modes of action.

The key of resistance management strategies is the reduction of selection pressure. This can be achieved by good agricultural practice which leads to less infection pressure (e.g. crop & stubble hygiene, use of less susceptible varieties, crop rotation, use of healthy seeds, etc).

Risk assessments for the target diseases on oilseed rape indicate a low or medium risk of resistance development. To minimise any possible risk of resistance occurring, the number of applications of Filan should be restricted to two per crop. Filan should be applied in a preventive manner in mixture with a non-cross resistant fungicide with activity against the target disease at a dose that will give robust control and as an integral part of a spray programme including fungicides with different modes of action.

For further advice on resistance management in SDHIs contact your agronomist or specialist advisor, and visit the FRAG-UK website

3. Crops

Oilseed rape

Time of Application

Apply Filan as a protectant spray. Two applications can be made from the four true leaf stage of the crop in the autumn, up to before 60% of pods have reached final size (GS 76).

Applications against *Phoma* should be made from early autumn at the first signs of *Phoma* leaf spot infection.

Applications against *Sclerotinia* should be made in high disease risk situations at early to full flower (GS 63-65 BBCH), before disease symptoms are visible.

Applications against *Alternaria* should be made from full flowering (GS 65 BBCH) up to before 60% of pods have reached final size (GS 76). Applications made after the end of flowering (GS69) may result in reduced levels of control.

4. Mixing and Application

4.1 Mixing

Never prepare more spray solution than is required.

Half fill the spray tank with clean water and start the agitation. Slowly add the required amount of Filan whilst re-circulating.

Rinse empty containers thoroughly, using an integrated pressure rinsing device or by manually rinsing three times. Add washings to tank at time of filling and dispose of container safely.

Add the remainder of the water and continue agitation until spraying is complete.

4.2 Application

Apply 0.5kg of Filan in 220 - 400 litres of water per hectare as a MEDIUM spray, as defined by BCPC. In dense crops use the higher volume to achieve adequate penetration and good coverage of the crop.

Where possible use narrow-wheeled high clearance application equipment. To ensure optimum spray coverage and minimize spray drift, adjust the spray boom to the appropriate height above the crop, in according to guidance provided by the sprayer and/or nozzle manufacture.

The following does not form part of the product label under S.I. No. 159 of 2012:

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of the Unfair Contract Terms Act 1977 or any similar applicable law.

Additional Product Safety Information

The information on this label is based on the best available information including data from test results

Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



http://www.agricentre.basf.ie/go/filan_sds

Alternatively, contact your supplier.

SPECIMEN -
2016 to date

Filan[®]

Filan is a protectant and systemic fungicide for use against early and late-season diseases in oilseed rape.

A water dispersible granule containing 50% w/w boscalid.

Risk and Safety Information

If medical advice is needed, have product container or label at hand.
Keep out of reach of children.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

PCS No.: 04325

To avoid risks to human health and the environment, comply with the instructions for use.

2.5 kg e

UN 3077
Packing Group III
Environmentally hazardous
substance, solid, N.O.S.
(contains boscalid 50%)
Marine Pollutant

Supplied by:
BASF Ireland Limited
P.O. Box 4, Earl Road
Cheadle Hulme, CHEADLE
Cheshire SK8 6QG, UK
Tel: 01 825 5701
Fax: 01 825 2038

Emergency Information
(24 hours freephone):
0049 180 227 3112
Technical Enquiries:
0044 (0)845 602 2553 (office hours)

Authorization holder:
BASF plc, P.O. Box 4,
Earl Road, Cheadle Hulme,
CHEADLE
Cheshire, SK8 6QG, UK


We create chemistry

® = Registered trademark of BASF

81107455IE1056