

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RESTRICTIONS OR WARNING

Only use on crops at risk of lodging.
This product must not be used on grass seed crops that will be grazed by livestock or cut for fodder.
Avoid spray drift onto neighbouring crops.
Take particular care to avoid overlapping spray swaths.
Do not apply if rain or frost is expected or if the crop is wet. Do not spray onto crops which are suffering stress, waterlogging, pest attack, disease or frost.

GENERAL INFORMATION

TRINEXIS is an emulsifiable concentrate formulation containing 250 g/L of trinexapac-ethyl. It is a foliar applied plant growth regulator (PGR) that can lead to lodging control. Treatment may lead to ears remaining erect through to harvest.

CROP-SPECIFIC INFORMATION

Crop	Rate of Use (L product/ha)	Timing
Winter wheat	0.4	Apply between leaf sheath erect stage and flag leaf fully emerged stage (GS30-39)
Spring wheat	0.4	Apply between leaf sheath erect stage and second node detectable stage (GS30-32)
Winter barley	0.4 or 0.6	Apply between leaf sheath erect stage and second node detectable stage (GS30-32) OR Apply between flag leaf just visible stage flag leaf fully emerged stage (GS37-39)
Spring barley	0.5	Apply between leaf sheath erect stage and second node detectable stage (GS30-32)
Winter and spring oats	0.4	Apply between leaf sheath erect stage and first node detectable stage (GS30-31)
Rye, triticale and Durum wheat	0.4	Apply between leaf sheath erect stage and second node detectable stage (GS30-32)
Ryegrass seed crops	0.8	Apply between leaf sheath erect stage and the first node detectable stage (GS30-31)

Spray Volume: For cereals and seed crops, apply TRINEXIS in 200-400 litres water/ha. The higher water volume may be used if the crop canopy is dense to aid penetration of the spray.

MIXING AND SPRAYING

Spray quality: To be applied using a conventional tractor mounted/trailed sprayer using a MEDIUM spray (as defined by the BCPC system).

Mixing and Application: Add the required quantity of TRINEXIS to a half-filled spray tank with continuous agitation. Add the remaining volume of water, maintaining agitation. Spray immediately after mixing and continue agitation until spraying is completed.
Wash out the sprayer and measuring equipment thoroughly with water and a wetting agent immediately after use. Finally wash out with water and drain.

COMPANY ADVISORY INFORMATION

This section is not part of the (EC) 1107/2009 Regulation. It provides additional advice on product use at the discretion of the Authorisation holder.

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling mixing or use or the weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

TRINEXIS

Contains 250 g/L trinexapac-ethyl as an emulsifiable concentrate (EC)

Plant growth regulator for plant height reduction, which can lead to lodging control in cereals and in ryegrass crops grown for seed production.

Important information

**FOR PROFESSIONAL USE ONLY.
FOR USE AS AN AGRICULTURAL PLANT GROWTH REGULATOR**

Crops	Maximum individual dose (L product/ha)	Maximum dose of application (L product/ha)	Max. total dose (L product/ha)	Latest time of application
Winter wheat	0.4	-	0.4	Before flag leaf sheath extending stage (GS41)
Spring wheat	0.4	-	0.4	Before third node detectable stage (GS33)
Winter barley	0.6	-	0.6	Before flag leaf sheath extending stage (GS41)
Spring barley	0.5	-	0.5	Before third node detectable stage (GS33)
Winter & spring oats	0.4	-	0.4	Before second node detectable stage (GS32)
Durum wheat, rye, triticale	0.4	-	0.4	Before third node detectable stage (GS33)
Grassland (seed crops)	0.8	-	0.8	Before second node detectable stage (GS32)

Other specific restrictions: This product must not be used on grass seed crops that will be grazed by livestock or cut for fodder. Treated grassland must not be grazed or cut for fodder.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY OFFENSE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

Net Contents: 1 litre e

Manufacturing date and Batch No.: see packaging

Manufacturer, Marketing Company and Approval holder:

Arysta LifeScience Benelux sprl

Rue de Renory 26/1

B-4102 Ougrée

Belgium

Tel. +32 4385 9711

24 hour emergency telephone number: +44 (0) 1235 239670



Risk & Safety Information

Warning

**May cause an allergic skin reaction.
Toxic to aquatic life with long lasting effects.**

Keep out of reach of children.

Do not eat, drink or smoke when using this product.

Wear protective gloves/protective clothing.

IF ON SKIN: Wash with plenty of soap and water.

Collect spillage.

Dispose of contents/container to a licensed waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.
PCS No. 05145

SAFETY PRECAUTIONS

Operator protection

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate.
WASH CONCENTRATE from skin or eyes immediately.
WASH HANDS before meals and after work.

Environmental protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Storage and disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.
WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.
DO NOT RE-USE CONTAINER FOR ANY PURPOSE.

PROTECT FROM FROST.

TRINEXIS