



TRIANUM-P

This product contains 1 % w/w *Trichoderma harzianum* Rifal strain T-22 spores (1 x 10⁸ spores/g) and 99 % w/w inert ingredients formulated as a water dispersible granule (WG).

Trianium-P is a biofungicide for the reduction of soil-borne *Pythium*, *Rhizoctonia* and *Fusarium* in protected edible and ornamental crops. It can be applied via drench application to the growing medium, or through drip-irrigation systems, or by spraying at sowing.

MANUFACTURED FOR

Koppert UK Ltd
8 Tudor Rose Court
53 Hollands Road
Haverhill
Suffolk, CB9 8PJ
Tel. 01440 704488

BATCH NO.
see top

USE BY
see top

NET CONTENT
500 g in PE lined cardboard container

PRODUCT NUMBER
11412

APPROVAL HOLDER AND MARKETING COMPANY
Koppert BV
PO Box 155
2650 Berkel en Rodenrijs
The Netherlands

COMPANY ADVISORY INFORMATION
This section is not part of the (EC) 1107/2009 Regulation.

SAFETY INFORMATION

- Read label before use
- Keep away from food, drink and animal feeding stuffs
- Keep out of the reach of children
- Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste
- Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.
- Keep only in the original container
- Wash hands and exposed skin before meals and after work
- Do not breathe dust
- Avoid contact with eyes
- In case of contact with eyes, rinse immediately with plenty of water and seek medical advice
- If medical advice is needed, have product container or label at hand
- Wear suitable protective clothing (coveralls), suitable protective gloves and suitable respiratory protective equipment* when handling or applying the product.

*Disposable filtering facepiece respirator to at least EN149 FFP3 or equivalent

Micro-organisms may have the potential to provoke sensitising reactions

OPERATOR PROTECTION

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:
Operators must wear suitable protective clothing (coveralls), suitable protective gloves and suitable respiratory protective equipment* when handling the product or applying the product.

*Disposable filtering facepiece respirator to at least EN149 FFP3 or equivalent.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS NO. 05486

TRIANUM-P



IMPORTANT INFORMATION
FOR USE ONLY AS AN AGRICULTURAL/HORTICULTURAL FUNGICIDE

User: Professional

Crop/situation Area of use	Maximum Individual dose	Maximum number of treatments	Maximum total dose	Last time of application
Protected cucumber, tomato, pepper, chilli and aubergine	15 kg/ha	2 per crop	30 kg/ha	At propagation
	and			
	15 g per 1000 plants at cultivation	12 per year	-	-
Protected ornamental plant production	15 kg/ha	2 per crop	30 kg/ha	At propagation
	and			
	15 g per 1000 plants at cultivation	4 per year	-	-
Protected lettuce and baby leaf production	15 kg/ha	1 per crop	15 kg/ha	-

Specific pest controlled:
Reduction of soil-borne *Pythium*, *Rhizoctonia* and *Fusarium*

Other Specific Restrictions:
A minimum interval of 14 days must be observed between applications at propagation
A minimum interval of 4 weeks must be observed between applications at cultivation



TRIANUM-P

APPLICATION AND DOSE

Method of application: drench, drip irrigation, glasshouse sprayer (at sowing).

Trianium-P should be used preventatively. For the best results, always use Trianium-P from propagation onwards, before occurrence of disease.

Use sufficient suspension to guarantee good penetration of the growing medium, but to maximise spore attachment to the plant roots, avoid run-off and excessive drainage for the first two days following application.

Do not use disinfectants in drip irrigation systems for at least one day before and two days after the application of Trianium-P. Introduce Trianium-P into the irrigation system AFTER the filter(s).

Trianium-P is not a systemic product, and is therefore not taken up by the roots.

Propagation fruiting vegetables and ornamentals
Apply 1.5 g/m² of cultivated area, suspended in 2.5-5 litres of water, immediately after sowing seeds, or sticking cuttings.

Propagation of lettuce and baby leaves
Apply 1.5 g/m² of cultivated area, suspended in 1.5 - 2 litres of water, immediately after sowing.

Cultivation after propagation
Apply 15 g/1000 plants directly after planting.
Use a volume of water equivalent to 10% of the substrate volume, or 100 litres/1000 plants.

Repeat the dose every 10 weeks, or after potting-on.

Disease reduction

The beneficial fungus *Trichoderma harzianum* T-22 out-competes plant pathogenic fungi for space and nutrients, colonising the plant roots ahead of the pathogens. It also has some activity as a mycoparasite by producing enzymes which break down the hyphae of the plant pathogenic fungi.

STORAGE

Trianium-P is a biological product and has a guaranteed shelf-life of six months at 4°C. The product should be stored at 4 – 8°C and used within six months of the date of manufacture.

Do not keep in domestic refrigerators

Do not freeze

KOPPERT BV guarantees that Trianium-P is fit for the purpose and conforms to the specifications stated on this label. KOPPERT BV gives no warranty, express or implied, including the warranties of commercially acceptable quality and/or fitness for any particular purpose concerning this material, other than those which are stated on this label, and will not be liable for any direct or indirect damage which results from incorrect use of the product or non-compliance with the directions of this label.

SPECIMEN -
2016 to date

TRIANUM-P

PCS NO. 05436

This product contains 1 % w/w *Trichoderma harzianum* Rifai strain T-22 spores (1 x 10⁹ spores/g) and 99 % w/w inert ingredients formulated as a water dispersible granule (WG).

Trianium-P is a biofungicide for the reduction of soil-borne *Pythium*, *Rhizoctonia* and *Fusarium* in protected edible and ornamental crops. It can be applied via drench application to the growing medium, or through drip-irrigation systems, or by spraying at sowing.

MANUFACTURED FOR

Koppert UK Ltd
8 Tudor Rose Court
53 Hollands Road
Haverhill
Suffolk, CB9 8PJ
Tel. 01440 704488

BATCH NO

see top

USE BY

see top

NET CONTENT

500 g in PE lined cardboard container

PRODUCT NUMBER

11412

APPROVAL HOLDER AND MARKETING COMPANY

Koppert BV
PO Box 155
2650 Berkel en Rodenrijs
The Netherlands.

COMPANY ADVISORY INFORMATION

This section is not part of the (EC) 1107/2009 Regulation.

**IMPORTANT INFORMATION
FOR USE ONLY AS AN AGRICULTURAL/HORTICULTURAL FUNGICIDE**

User: Professional

Crop/situation Area of use	Maximum Individual dose	Maximum number of treatments	Maximum total dose	Last time of application
Protected cucumber, tomato, pepper, chilli and aubergine	15 kg/ha	2 per crop	30 kg/ha	At propagation
	and			
	15 g per 1000 plants at cultivation	12 per year	-	-
Protected ornamental plant production	15 kg/ha	2 per crop	30 kg/ha	At propagation
	and			
	15 g per 1000 plants at cultivation	4 per year	-	-
Protected lettuce and baby leaf production	15 kg/ha	1 per crop	15 kg/ha	-

Specific pest controlled:
Reduction of soil-borne Pythium, Rhizoctonia and Fusarium

Other Specific Restrictions:
A minimum interval of 14 days must be observed between applications at propagation
A minimum interval of 4 weeks must be observed between applications at cultivation

Product to be applied at temperatures above 10° C only

READ ALL OTHER SAFETY PRECAUTIONS AND DIRECTIONS FOR USE BEFORE USE

SAFETY INFORMATION

Read label before use
Keep away from food, drink and animal feeding stuffs
Keep out of the reach of children
Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste
Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.
Keep only in the original container
Wash hands and exposed skin before meals and after work
Do not breathe dust
Avoid contact with eyes
In case of contact with eyes, rinse immediately with plenty of water and seek medical advice
If medical advice is needed, have product container or label at hand
Wear suitable protective clothing (coveralls), suitable protective gloves and suitable respiratory protective equipment* when handling or applying the product. *Disposable filtering facepiece respirator to at least EN149 FFP3 or equivalent

Micro-organisms may have the potential to provoke sensitising reactions

OPERATOR PROTECTION

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

Operators must wear suitable protective clothing (coveralls), suitable protective gloves and suitable respiratory protective equipment* when handling the product or applying the product.
*Disposable filtering facepiece respirator to at least EN149 FFP3 or equivalent.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No. 05486

DIRECTIONS FOR USE

GENERAL INFORMATION

Trianum-P protects plants against infection by soil-borne pathogens via the root system. This can enhance the growth and development of roots and aerial parts of the plant. Trianum-P can be used under various environmental conditions. It grows under a range of temperatures above 10° C (maximum temperature 34° C), at a pH between 4-8, in many types of growing media, and on the roots of a wide range of plants.

PREPARATION OF SUSPENSION

Suspend the required dose of Trianum-P in water, as 1 volume of Trianum-P in 5 volumes of water.
Mix thoroughly.
Add suspension to required amount of water to obtain final drench volume (see below).
Agitate continuously to maintain an even distribution of Trianum-P spores, as they can precipitate out.
Apply the suspension on the day of preparation.

No PHI (minimum pre-harvest interval) is needed between treatment with Trianum-P and harvest.

COMPATIBILITY

Trianum-P should not be applied in mixture with other pesticides or liquid fertilisers. The effect of other pesticides applied either before or after Trianum-P on its effectiveness has not been fully established. Some pesticides, particularly fungicides, may reduce effectiveness of Trianum-P.

It is compatible with natural enemies, and can be used in integrated crop management systems.

Trianum-P has been used on a wide range of crops under a wide range of conditions and no phytotoxicity has been observed.

Application equipment should be cleaned thoroughly after use.

Consult processor before using on crops intended for processing.